

**Use identities to simplify each expression:**

$$1. \quad \frac{\tan x \csc x}{\sec x} = \frac{\left(\frac{\sin x}{\cos x}\right)\left(\frac{1}{\sin x}\right)}{\frac{1}{\cos x}} = \frac{\frac{1}{\cos x}}{\frac{1}{\cos x}} = \left(\frac{1}{\cos x}\right)\left(\frac{\cos x}{1}\right) = 1$$

$$2. \quad \tan^2 x - \frac{\sin(-x)}{\sin x} = \tan^2 x - \left(\frac{-\sin x}{\sin x}\right) = \tan^2 x + 1 = \sec^2 x$$

**Multiply and simplify:**

$$3. \quad \sin \theta \cos \theta (\tan \theta + \cot \theta) = \sin \theta \cos \theta \left( \frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} \right) = \sin^2 \theta + \cos^2 \theta = 1$$

**Factor and simplify:**

$$4. \quad \sin^2 x \tan^2 x + \sin^2 x = \sin^2 x (\tan^2 x + 1) = \sin^2 x (\sec^2 x) = \sin^2 x \left( \frac{1}{\cos^2 x} \right) = \tan^2 x$$

$$5. \quad \frac{\sin x \cos x}{\tan x} = 1 - \sin^2 x$$

$$\begin{aligned} \frac{\sin x \cos x}{\tan x} &= \frac{\sin x \cos x}{\frac{\sin x}{\cos x}} \\ &= \sin x \cos x \left( \frac{\cos x}{\sin x} \right) \\ &= \cos^2 x \\ &= 1 - \sin^2 x \end{aligned}$$

$$6. \quad \cot(-x) = \frac{1 - \sin^2 x}{\cos(-x) \sin(-x)}$$

$$\begin{aligned} \cot(-x) &= \frac{1 - \sin^2 x}{\cos(-x) \sin(-x)} = \frac{\cos^2 x}{\cos x (-\sin x)} \\ &= \frac{\cos x}{-\sin x} \\ &= -\cot x \end{aligned}$$

$$7. \quad \frac{\sin 2\beta}{2 \csc \beta} = \sin^2 \beta \cos \beta$$

$$\begin{aligned} \frac{\sin 2\beta}{2 \csc \beta} &= \frac{2 \sin \beta \cos \beta}{\frac{2}{\sin \beta}} \\ &= 2 \sin \beta \cos \beta \left( \frac{\sin \beta}{2} \right) \\ &= \sin^2 \beta \cos \beta \end{aligned}$$

$$8. \frac{1}{\sec \theta - 1} - \frac{1}{\sec \theta + 1} = 2 \cot^2 \theta$$

$$\begin{aligned} \frac{1}{\sec \theta - 1} - \frac{1}{\sec \theta + 1} &= \left( \frac{1}{\sec \theta - 1} \right) \left( \frac{\sec \theta + 1}{\sec \theta + 1} \right) - \left( \frac{1}{\sec \theta + 1} \right) \left( \frac{\sec \theta - 1}{\sec \theta - 1} \right) \\ &= \frac{\sec \theta + 1 - (\sec \theta - 1)}{(\sec \theta + 1)(\sec \theta - 1)} \\ &= \frac{\sec \theta + 1 - \sec \theta + 1}{\sec^2 \theta - 1} \\ &= \frac{2}{\tan^2 \theta} \\ &= 2 \cot^2 \theta \end{aligned}$$

$$9. \cos(3x) = \cos x(1 - 4 \sin^2 x)$$

$$\begin{aligned} \cos(3x) &= \cos(2x + x) \\ &= \cos 2x \cos x - \sin 2x \sin x \\ &= (1 - 2 \sin^2 x) \cos x - (2 \sin x \cos x) \sin x \\ &= \cos x - 2 \sin^2 x \cos x - 2 \sin^2 x \cos x \\ &= \cos x - 4 \sin^2 x \cos x \\ &= \cos x(1 - 4 \sin^2 x) \end{aligned}$$

$$10. \sin^2 \left( \frac{x}{2} \right) = \frac{\csc^2 x - \cot^2 x}{2 \csc^2 x + 2 \csc x \cot x}$$

$$\begin{aligned} \sin^2 \left( \frac{x}{2} \right) &= \left( \pm \sqrt{\frac{1 - \cos x}{2}} \right)^2 \\ &= \frac{1 - \cos x}{2} \end{aligned}$$

$$\begin{aligned} \frac{\csc^2 x - \cot^2 x}{2 \csc^2 x + 2 \csc x \cot x} &= \frac{1}{\frac{2}{\sin^2 x} + \left( \frac{2}{\sin x} \right) \left( \frac{\cos x}{\sin x} \right)} \\ &= \frac{1}{\frac{2 + 2 \cos x}{\sin^2 x}} \\ &= \frac{\sin^2 x}{2 + 2 \cos x} \\ &= \frac{1 - \cos^2 x}{2 + 2 \cos x} \\ &= \frac{(1 - \cos x)(1 + \cos x)}{2(1 + \cos x)} \\ &= \frac{1 - \cos x}{2} \end{aligned}$$

11. Use trigonometric identities to find the values of the other five trigonometric functions if  $\cos \alpha = 1/\sqrt{5}$  and  $\alpha$  is in quadrant IV.

In quadrant IV, cosine and secant are positive, and all the other functions are negative.

$$\sin^2 \alpha + \cos^2 \alpha = 1$$

$$\csc \alpha = \frac{1}{\sin \alpha} = \frac{1}{-2/\sqrt{5}} = -\frac{\sqrt{5}}{2}$$

$$\sin^2 \alpha + \left(\frac{1}{\sqrt{5}}\right)^2 = 1$$

$$\tan \alpha = \frac{\sin \alpha}{\cos \alpha} = \frac{-2/\sqrt{5}}{1/\sqrt{5}} = -2$$

$$\sec \alpha = \frac{1}{\cos \alpha} = \frac{1}{1/\sqrt{5}} = \sqrt{5}$$

$$\sin^2 \alpha + \frac{1}{5} = 1$$

$$\cot \alpha = \frac{1}{\tan \alpha} = -\frac{1}{2}$$

$$\sin^2 \alpha = \frac{4}{5}$$

$$\sin \alpha = -\sqrt{\frac{4}{5}} = -\frac{2}{\sqrt{5}}$$

**Find the exact value by using a sum or difference identity:**

$$\begin{aligned} 12. \quad \cos\left(\frac{7\pi}{12}\right) &= \cos\left(\frac{\pi}{3} + \frac{\pi}{4}\right) \\ &= \cos\left(\frac{\pi}{3}\right)\cos\left(\frac{\pi}{4}\right) - \sin\left(\frac{\pi}{3}\right)\sin\left(\frac{\pi}{4}\right) \\ &= \left(\frac{1}{2}\right)\left(\frac{\sqrt{2}}{2}\right) - \left(\frac{\sqrt{3}}{2}\right)\left(\frac{\sqrt{2}}{2}\right) = \frac{\sqrt{2}}{4} - \frac{\sqrt{6}}{4} = \frac{\sqrt{2} - \sqrt{6}}{4} \end{aligned}$$

$$\begin{aligned} 13. \quad \sin\left(\frac{\pi}{12}\right) &= \sin\left(\frac{\pi}{3} - \frac{\pi}{4}\right) \\ &= \sin\left(\frac{\pi}{3}\right)\cos\left(\frac{\pi}{4}\right) - \cos\left(\frac{\pi}{3}\right)\sin\left(\frac{\pi}{4}\right) \\ &= \left(\frac{\sqrt{3}}{2}\right)\left(\frac{\sqrt{2}}{2}\right) - \left(\frac{1}{2}\right)\left(\frac{\sqrt{2}}{2}\right) = \frac{\sqrt{6}}{4} - \frac{\sqrt{2}}{4} = \frac{\sqrt{6} - \sqrt{2}}{4} \end{aligned}$$


$$14. \quad \tan(165^\circ) = \tan(120^\circ + 45^\circ) = \frac{\tan(120^\circ) + \tan(45^\circ)}{1 - \tan(120^\circ)\tan(45^\circ)} = \frac{-\sqrt{3} + 1}{1 - (-\sqrt{3})(1)} = \frac{-\sqrt{3} + 1}{1 + \sqrt{3}}$$

**Use the sum/difference identities to simplify each expression.**

$$15. \quad \cos 75^\circ \cos 60^\circ - \sin 75^\circ \sin 60^\circ = \cos(75^\circ + 60^\circ) = \cos 135^\circ = -\sqrt{2}/2$$


$$16. \quad \sin 80^\circ \cos 50^\circ - \cos 80^\circ \sin 50^\circ = \sin(80^\circ - 50^\circ) = \sin 30^\circ = 1/2$$

17. Find  $\sin(A+B)$  if  $\sin A = -5/13$  and  $\cos B = 2/5$ , with  $A$  in quadrant III and  $B$  in quadrant I.



$$x = -\sqrt{13^2 - (-5)^2} = -12$$

$$\cos A = -\frac{12}{13}$$



$$y = \sqrt{5^2 - 2^2} = \sqrt{21}$$


$$\sin B = \frac{\sqrt{21}}{5}$$

$$\sin(A+B) = \sin A \cos B + \cos A \sin B$$

$$= \left(-\frac{5}{13}\right)\left(\frac{2}{5}\right) + \left(-\frac{12}{13}\right)\left(\frac{\sqrt{21}}{5}\right)$$


$$= \frac{-10 - 12\sqrt{21}}{65}$$

18. Find  $\cos(\alpha - \beta)$  if  $\sin \alpha = 15/17$  and  $\sin(\beta) = -1/3$ , with  $\alpha$  in quadrant II and  $\beta$  in quadrant IV.



$$x = -\sqrt{17^2 - 15^2} = -8$$

$$\cos \alpha = -\frac{8}{17}$$



$$x = \sqrt{3^2 - (-1)^2} = \sqrt{8} = 2\sqrt{2}$$

$$\cos \beta = \frac{2\sqrt{2}}{3}$$

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$= \left(-\frac{8}{17}\right)\left(\frac{2\sqrt{2}}{3}\right) + \left(\frac{15}{17}\right)\left(-\frac{1}{3}\right)$$

$$= \frac{-16\sqrt{2} - 15}{51}$$

**Find the exact value by using a half-angle identity.**

19. We know  $\sin(-\pi/8)$  is negative because  $-\pi/8$  is in quadrant IV.

$$\sin(-\pi/8) = \sin\left(\frac{-\pi/4}{2}\right) = -\sqrt{\frac{1 - \cos(-\pi/4)}{2}} = -\sqrt{\left(\frac{1 - \sqrt{2}/2}{2}\right)\left(\frac{2}{2}\right)} = -\sqrt{\frac{2 - \sqrt{2}}{4}} = \frac{-\sqrt{2 - \sqrt{2}}}{2}$$

20. We know that  $\tan(3\pi/8)$  is positive because  $3\pi/8$  is in quadrant I.

$$\begin{aligned} \tan(3\pi/8) &= \tan\left(\frac{3\pi/4}{2}\right) \\ &= \frac{1 - \cos(3\pi/4)}{\sin(3\pi/4)} = \frac{1 - \left(-\frac{\sqrt{2}}{2}\right)}{\frac{\sqrt{2}}{2}} = \left(\frac{1 + \frac{\sqrt{2}}{2}}{\frac{\sqrt{2}}{2}}\right) \cdot \frac{2}{2} = \frac{2 + \sqrt{2}}{\sqrt{2}} \\ &= \left(\frac{2 + \sqrt{2}}{\sqrt{2}}\right)\left(\frac{\sqrt{2}}{\sqrt{2}}\right) = \frac{2\sqrt{2} + 2}{2} = \sqrt{2} + 1 \end{aligned}$$


**Use the given information to find the exact value of the trigonometric function.**

21. Find  $\cos\left(\frac{\alpha}{2}\right)$  if  $\cos \alpha = \frac{1}{4}$ , and  $\alpha$  is in Quadrant IV.

$270^\circ < \alpha < 360^\circ \Rightarrow 135^\circ < \alpha/2 < 180^\circ$ , so  $\alpha/2$  is in quadrant II and  $\cos(\alpha/2)$  is negative.

$$\cos\left(\frac{\alpha}{2}\right) = -\sqrt{\frac{1 + \cos \alpha}{2}} = -\sqrt{\frac{1 + 1/4}{2}} = -\sqrt{\frac{5/4}{2}} = -\sqrt{\frac{5}{8}} = -\frac{\sqrt{5}}{2\sqrt{2}}$$

22. Find  $\sin(2\theta)$  if  $\sin(\theta) = \frac{5}{13}$ , and  $\theta$  is in quadrant II.



$$\sin(2\theta) = 2 \sin \theta \cos \theta = 2 \left( \frac{5}{13} \right) \left( -\frac{12}{13} \right) = -\frac{120}{169}$$

$$x = -\sqrt{13^2 - 5^2} = -12$$

$$\cos \theta = -\frac{12}{13}$$

**Find the exact value without using a calculator.**

23.  $\cos^{-1}\left(-\frac{1}{2}\right) = \frac{2\pi}{3}$

Remember,  $\cos^{-1}$  is between 0 and  $\pi$ .

24.  $\arctan(1) = \frac{\pi}{4}$

Remember,  $\arctan$  is between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$ .

25.  $\sec^{-1}\left(\frac{2\sqrt{3}}{3}\right) = \sec^{-1}\left(\frac{2}{\sqrt{3}}\right) = \cos^{-1}\left(\frac{\sqrt{3}}{2}\right) = \frac{\pi}{6}$

Remember,  $\sec^{-1}$  is between 0 and  $\pi$ .

**Find the exact value in degrees without using a calculator.**

26.  $\csc^{-1}(-\sqrt{2}) = \sin^{-1}\left(-\frac{1}{\sqrt{2}}\right) = -45^\circ$

Remember,  $\csc^{-1}$  is between  $-90^\circ$  and  $90^\circ$ .

27.  $\arcsin\left(\frac{1}{2}\right) = 30^\circ$

Remember,  $\arcsin$  is between  $-90^\circ$  and  $90^\circ$ .

28.  $\cot^{-1}(-\sqrt{3}) = \cot^{-1}\left(\frac{-\sqrt{3}/2}{1/2}\right) = 150^\circ$

Remember,  $\cot^{-1}$  is between  $0^\circ$  and  $180^\circ$ .

$$\left( \cot x = \frac{\cos x}{\sin x}, \text{ so } \cos x = -\frac{\sqrt{3}}{2} \text{ and } \sin x = \frac{1}{2}. \right)$$

**Find the approximate value using a calculator.**

29.  $\sec^{-1}(-1.5643) = \cos^{-1}\left(\frac{1}{-1.5643}\right) = 2.26$

**Find all values for  $x$  in the interval  $[0, 2\pi]$  that satisfy the equation.**

30.  $\tan x = \sqrt{3}$


$$x = \frac{\pi}{3} + k\pi$$

$$\left\{ \frac{\pi}{3}, \frac{4\pi}{3} \right\}$$

**Find the exact value of each composition.**

$$31. \csc \left( \underbrace{\cos^{-1} \left( \frac{5}{13} \right)}_{\theta} \right)$$

What is the cosecant of the angle whose cosine is  $\frac{5}{13}$ ?




$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{5}{13}$$

$$\text{opp} = \sqrt{13^2 - 5^2} = 12$$

$$\csc \theta = \frac{\text{hyp}}{\text{opp}} = \frac{13}{12}$$

$$32. \cos \left( \underbrace{\arctan \left( \frac{8}{15} \right)}_{\theta} \right)$$

What is the cosine of the angle whose tangent is  $\frac{8}{15}$ ?



$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{8}{15}$$


$$\text{hyp} = \sqrt{8^2 + 15^2} = 17$$

$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{15}{17}$$

**Find an equivalent algebraic expression for each composition.**

$$33. \tan \left( \underbrace{\arccos(x)}_{\theta} \right)$$

What is the tangent of the angle whose cosine is  $x$ ?




$$\cos \theta = \frac{\text{adj}}{\text{hyp}} = \frac{x}{1}$$

$$\text{opp} = \sqrt{1^2 - x^2} = \sqrt{1 - x^2}$$

$$\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{\sqrt{1 - x^2}}{x}$$

$$34. \sec \left( \underbrace{\arcsin(x)}_{\theta} \right)$$

What is the secant of the angle whose sine is  $x$ ?



$$\sin \theta = \frac{\text{opp}}{\text{hyp}} = \frac{x}{1}$$

$$\text{adj} = \sqrt{1^2 - x^2} = \sqrt{1 - x^2}$$

$$\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{1}{\sqrt{1 - x^2}}$$

**Find the acute angle  $\theta$ , to the nearest hundredth of a degree for the given function value.**

$$35. \csc \theta = 6.354 \Rightarrow \sin \theta = \frac{1}{6.354}$$

$$\theta = \sin^{-1} \left( \frac{1}{6.354} \right) = 9.05^\circ$$

$$36. \sec \theta = 4.321 \Rightarrow \cos \theta = \frac{1}{4.321}$$

$$\theta = \cos^{-1} \left( \frac{1}{4.321} \right) = 76.62^\circ$$

**Find all real numbers that satisfy the equation.**

37.  $2 \cos x + \sqrt{3} = 0$

$$2 \cos x = -\sqrt{3}$$

$$\cos x = -\frac{\sqrt{3}}{2}$$

$$\left\{ x \mid x = \frac{5\pi}{6} + 2k\pi \text{ or } x = \frac{7\pi}{6} + 2k\pi \right\}$$

**Find all real numbers in the interval  $[0, 2\pi)$  that satisfy the equation.**

38.  $\sqrt{2} \sin(3x) = 1$

$$\sin(3x) = \frac{1}{\sqrt{2}}$$

$$3x = \frac{\pi}{4} + 2k\pi \text{ or } 3x = \frac{3\pi}{4} + 2k\pi$$

$$x = \frac{\pi}{12} + \frac{2k\pi}{3} \text{ or } x = \frac{\pi}{4} + \frac{2k\pi}{3}$$

$$\frac{\pi}{12} + \frac{2\pi}{3} = \frac{\pi}{12} + \frac{8\pi}{12} = \frac{9\pi}{12} = \frac{3\pi}{4}$$

$$\frac{\pi}{4} + \frac{2\pi}{3} = \frac{3\pi}{12} + \frac{8\pi}{12} = \frac{11\pi}{12}$$

$$\left\{ \frac{\pi}{12}, \frac{3\pi}{4}, \frac{\pi}{4}, \frac{11\pi}{12} \right\}$$

**Find all real numbers that satisfy the equation.**

39.  $2 \cos^2 x = \cos x$

$$2 \cos^2 x - \cos x = 0$$

$$\cos x(2 \cos x - 1) = 0$$

$$\cos x = 0 \text{ or } 2 \cos x - 1 = 0$$

$$\cos x = 0 \quad \text{or} \quad \cos x = \frac{1}{2}$$

$$\frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \text{ etc.} \quad \frac{\pi}{3}, \frac{5\pi}{3}$$

$$\left\{ x \mid x = \frac{\pi}{2} + k\pi \text{ or } x = \frac{\pi}{3} + 2k\pi \text{ or } x = \frac{5\pi}{3} + 2k\pi \right\}$$

**Find all angles in the interval  $[0^\circ, 360^\circ)$  that satisfy the equation.**

40.  $2 \sin x \cos x = 1$

$$\sin(2x) = 1$$

$$2x = 90^\circ + 360^\circ k$$

$$x = 45^\circ + 180^\circ k$$

$$\{45^\circ, 225^\circ\}$$